

THE CHRONIC TOXICITY OF AMMONIA, NITRITE AND NITRATE ON JUVENILE *Farfantepenaeus brasiliensis* (CRUSTACEA: DECAPODA)*

Bruno Ribeiro de CAMPOS¹; Plínio Schmidt FURTADO¹; Fernando D'INCAO¹; Luis POERSCH¹; Wilson WASIELESKY¹

ABSTRACT

In general, the adverse effect of a chemical compound present in water varies with the concentration and time of exposure to the compound, the nature of the chemical species and age of the exposed organisms. Thus, nitrogen does not necessarily cause adverse effects on shrimp, but may, instead, promote sub-lethal effects by long-term exposure. Juvenile *Farfantepenaeus brasiliensis* (initial mean weight = 0.61 g \pm 0.07) were exposed to sub-lethal concentrations of ammonia (0.44 and 0.88 mg L⁻¹), nitrite (5.30 and 10.60 mg L⁻¹) and nitrate (45.60 and 91.20 mg L⁻¹) corresponding to the safe levels for the species. After 40 days of exposure of juveniles to ammonia, nitrite and nitrate, all groups differed significantly ($p < 0.05$) from the control group regarding the growth and survival. Based on the results, it was determined that the shrimp *F. brasiliensis* was susceptible to nitrogen compounds in concentrations equivalent to supposedly safe levels previously proposed for the specie. Thus, the security levels of ammonia, nitrite and nitrate for pink shrimp juveniles were 0.88 mg L⁻¹, 10.60 mg L⁻¹ and 91.20 mg L⁻¹, respectively.

Keywords: nitrogen compounds; performance; pink shrimp

TOXICIDADE CRÔNICA DA AMÔNIA, NITRITO E NITRATO EM JUVENIS DE *Farfantepenaeus brasiliensis* (CRUSTACEA: DECAPODA)

RESUMO

Em geral, o efeito adverso de um composto químico presente na água varia com a concentração, o tempo de exposição ao composto, à natureza do produto químico e a idade das espécies de organismos expostos. Assim, o nitrogênio não necessariamente causa efeitos adversos, mas pode, em vez disso, promover efeitos subletais por meio da exposição em longo prazo. Juvenis de *Farfantepenaeus brasiliensis* (peso médio inicial = 0,61 g \pm 0,07) foram expostos a concentrações subletais de amônia (0,44 e 0,88 mg L⁻¹), nitrito (5,30 e 10,60 mg L⁻¹) e nitrato (45,60 e 91,20 mg L⁻¹) correspondente aos "níveis de segurança" para a espécie. Após 40 dias de exposição dos juvenis à amônia, nitrito e nitrato, todos os grupos diferiram significativamente ($p < 0,05$) do grupo controle em relação ao crescimento e sobrevivência. Com base nos resultados, o camarão *F. brasiliensis* foi susceptível aos compostos nitrogenados em concentrações equivalentes aos níveis supostamente seguros anteriormente propostos para a espécie. Assim, os níveis de segurança de amônia, nitrito e nitrato propostos para juvenis de camarão-rosa são 0,88 mg L⁻¹, 10,60 mg L⁻¹ e 91,20 mg L⁻¹, respectivamente.

Palavras chave: compostos nitrogenados; desempenho zootécnico; camarão-rosa

Artigo Científico: Recebido em 26/03/2014 – Aprovado em 02/04/2015

¹ Programa de Pós-graduação em Aquicultura, Instituto de Oceanografia, Universidade Federal do Rio Grande (FURG). Avenida Itália, km 8 – CEP: 96201-900 – Rio Grande – RS – Brasil. e-mail: brcampos@yahoo.com; pliniofs@yahoo.com.br; dincao@mikrus.com.br; lpoersch@mikrus.com.br; manow@mikrus.com.br (autor correspondente)

* Financial support: National Council for Scientific and Technological Development (CNPq), Ministry of Fishery and Aquaculture (MPA) and Coordination for the Improvement of Higher Level Personnel (CAPES)

INTRODUCTION

Knowledge of water quality parameters and their maintenance within tolerance limits for a species are essential requirements in any aquaculture system (KINNE, 1976). These factors decisively determine the success or failure of aquaculture (OSTRENSKY and WASIELESKY, 1995).

The nitrogenous waste from the excretion of farmed organisms and feed degradation frequently deteriorate the environment where these organisms are raised (TOMASSO, 1994). The nitrogen compounds can damage the gill tissues and affect oxygen consumption by the farmed organisms and/or cause their death (LIN and CHEN, 2003; KUHN *et al.*, 2010; BARBIERI, 2010; CAMPOS *et al.*, 2012; BARBIERI *et al.*, 2014).

The concentration and time required for a compound to produce an adverse effect vary according to the chemical agent and the type and severity of the effect. Adverse or toxic effects can be produced in the laboratory or in the natural environment through lethal (high concentrations for a short period of time) or chronic (sub-lethal concentrations over a long period of time) exposure to the chemical pollutant (RAND and PETROCELLI, 1985). The sensitivity of the organisms to a toxic substance also may vary according to their stage of development and their state of health (WAJSBROT *et al.*, 1993; COBO *et al.*, 2014).

In spite of the great majority of commercial farming that is performed with the exotic species *Litopenaeus vannamei*, some native species of marine shrimp, such as *Farfantepenaeus brasiliensis*, have already demonstrated the potential for culture (LOPES *et al.*, 2009). The application of

low-cost structures for culturing native shrimp (i.e., cages or pens in natural water bodies) allows the inclusion of low-income communities in this activity (WASIELESKY *et al.*, 2004). JENSEN (2012) recently concluded that *F. brasiliensis* is potentially suited for farming in bioflocs systems with a stocking density of up to 100 shrimp m⁻² during the nursery stage and up to 75 shrimp m⁻² during the grow-out stage.

The aim of this study was to evaluate the effects of sub-lethal concentrations of ammonia, nitrite and nitrate on the survival and growth of juvenile *F. brasiliensis* reared under laboratory conditions.

MATERIAL AND METHODS

Juveniles of *F. brasiliensis* were reared at the Marine Aquaculture Station hatchery, Federal University of Rio Grande (FURG), Rio Grande do Sul State, Brazil. The animals were kept in a 1,000 L tank with a controlled temperature and photoperiod (25 °C and 12L:12D, respectively), under constant aeration and with a salinity adjusted to 28.

The experiments were conducted in 200 L tanks, where thirty juveniles of *F. brasiliensis* (initial mean weight = 0.61 g ± 0.07), at a density of 120 shrimps m⁻³ were used per experimental unit in all treatments. The experiment consisted of a control group (without adding nitrogen) and six treatments, each with three replicates (21 tanks): two concentrations of ammonia (TAN), two of nitrite (NO₂⁻) and two of nitrate (NO₃⁻), which are presented in Table 1. These concentrations corresponded to the "safe levels" (SL) for this species (CAMPOS *et al.*, 2012) and half concentrations of those values (50%).

Table 1. Concentrations of ammonia, nitrite and nitrate used in the experiments.

Nitrogen compounds	Treatment	Concentration (mg L ⁻¹)
Without adding nitrogen	Control	0.0
		0.0
Ammonia	TAN-50%	0.44
	TAN-SL	0.88
Nitrite	NO ₂ ⁻ -50%	5.30
	NO ₂ ⁻ -SL	10.59
Nitrate	NO ₃ ⁻ -50%	45.60
	NO ₃ ⁻ -SL	91.20

The concentrations were obtained from stock solutions prepared with ammonium chloride p.a. (Synth®), sodium nitrite p.a. (Synth®) and sodium nitrate p.a. (Synth®).

The experiment lasted 40 days, and the water for each treatment was completely renewed every 48 h to maintain the desired concentrations. The excreta were siphoned daily and aeration was provided constantly. The water temperature was maintained at 25 °C using heaters with a thermostat (Visi-therm) and the salinity was kept at 28.

The water samples were collected daily from experimental units to measure salinity and pH with an optical refractometer (Atago, model 103) and a digital pH meter (DMpH-1, Digimed, precision 0.01), respectively. The concentrations of total ammonia (TAN) ($\text{NH}_3 + \text{NH}_4^+$), nitrite (NO_2^-) and nitrate (NO_3^-) were analyzed daily, according to the methodologies proposed by UNESCO (1983), BENDSCHNEIDER and ROBINSON (1952) and AMINOT and CHAUSSEPIED (1983), respectively. The oxygen concentrations and temperature were monitored daily with an oximeter (model OXI-315i, WTW).

The shrimp were fed twice a day (08:00 h and 16:00 h) with a commercial ration (\varnothing of pellets = 1.6 mm; 40% crude protein and 8% lipid) offered in trays at a proportion of 10% of the total biomass of each experimental unit. Every 10 days, 10

juveniles of each experimental unit were weighed with an electronic digital scale (± 0.01 g; SD-Marte®) to analyze the growth and feed adjustment. The mortality was verified every 24 h. Animals were considered dead when they were still and did not respond to mechanical stimuli with a glass cane (LIN and CHEN, 2003) and were immediately removed from the tank. The following performance parameters were determined: survival final (S; %) = $(\text{final } n / \text{initial } n) \times 100$ (where n = number of shrimp); final mean weight (g) = final biomass of live shrimp / total n ; specific growth rate (SGR; % day^{-1}) = $100 \times [(\ln \text{ final weight} - \ln \text{ initial weight}) / \text{days of experiment}]$.

The survival, wet weight and specific growth rate data were evaluated with an analysis of variance (one-way ANOVA) and validated in terms of the assumptions of the method (Levene Test and Kolmogorov-Smirnov). If significant differences were observed ($p < 0.05$), a Tukey test was applied to determine the moment when a significant toxic effect appeared. The percentage values were arcsine transformed (ZAR, 1996).

RESULTS

The mean values of pH, dissolved oxygen, temperature, salinity and nitrogen compounds registered for each treatment during the experimental period are presented in Table 2.

Table 2. Mean concentration (mg L^{-1}) values of ammonia, nitrite, nitrate, temperature (T; °C), dissolved oxygen (D.O.; mg L^{-1}), salinity and pH in the different treatments during the 40-day experiment. The data correspond to the mean \pm standard deviation.

Treatment	Concentration (mg L^{-1})	pH	D.O. (mg L^{-1})	T°C	Salinity
Control	-	8.31 \pm 0.13	6.32 \pm 0.24	24.96 \pm 0.21	28.03 \pm 0.50
TAN-50%	0.46 \pm 0.05	8.27 \pm 0.11	6.30 \pm 0.21	24.90 \pm 0.23	28.05 \pm 0.30
TAN-SL	0.88 \pm 0.08	8.32 \pm 0.15	6.32 \pm 0.21	24.87 \pm 0.26	28.00 \pm 0.50
NO_2^- -50%	5.32 \pm 0.18	8.27 \pm 0.13	6.36 \pm 0.30	24.86 \pm 0.27	28.05 \pm 0.50
NO_2^- -SL	10.59 \pm 0.25	8.31 \pm 0.16	6.23 \pm 0.27	24.88 \pm 0.22	28.10 \pm 0.40
NO_3^- -50%	45.61 \pm 0.58	8.30 \pm 0.15	6.40 \pm 0.22	24.96 \pm 0.24	28.15 \pm 0.40
NO_3^- -SL	91.29 \pm 0.36	8.27 \pm 0.16	6.36 \pm 0.20	24.95 \pm 0.23	28.05 \pm 0.53

The specific growth rate was significantly higher ($p < 0.05$) in the control when compared with the other treatments. Among the nitrogen treatments, there were no significant differences ($p > 0.05$). The specific growth and survival data

are shown in Table 3. The greatest final mean weight and percentage of survival was obtained in the control group and the lowest survival rate was observed with the nitrate treatment (NO_3^- -SL).

Table 3. Data (mean \pm standard deviation) for final mean weight (g), specific growth rate (SGR; % day⁻¹) and survival (%) of *Farfantepenaeus brasiliensis* juveniles after 40 days of exposure to ammonia (TAN-50% and TAN-SL), nitrite (NO₂-50% and NO₂-SL) and nitrate (NO₃-50% and NO₃-SL).

Treatment	Final Weight (g)	SGR (%/day)	Survival (%)
Control	1.11 \pm 0.27 ^a	1.53 \pm 0.15 ^a	77.78 \pm 3.85 ^a
TAN-50%	0.94 \pm 0.14 ^{ab}	1.02 \pm 0.20 ^b	64.44 \pm 3.85 ^{bc}
TAN-SL	0.90 \pm 0.03 ^b	1.01 \pm 0.09 ^b	60.00 \pm 6.67 ^{bc}
NO ₂ -50%	0.88 \pm 0.08 ^b	0.94 \pm 0.07 ^b	67.78 \pm 1.92 ^b
NO ₂ -SL	0.90 \pm 0.11 ^b	0.89 \pm 0.15 ^b	58.89 \pm 1.92 ^{bc}
NO ₃ -50%	0.94 \pm 0.05 ^b	0.93 \pm 0.03 ^b	67.78 \pm 1.92 ^b
NO ₃ -SL	0.92 \pm 0.04 ^b	1.00 \pm 0.04 ^b	56.67 \pm 3.33 ^c

Different superscript letters in the same column indicate significantly different means among the treatments ($p < 0.05$).

The highest growth rate of juvenile *F. brasiliensis* occurred in the control group when compared to the other treatments. Significant differences were found ($p < 0.05$) after 30 days in treatments TAN-SL, NO₂-50% and NO₂-SL when compared with the control treatment. On

day 40, significant differences ($p < 0.05$) were observed between the control group and the nitrogen treatments. The increase in the wet weight with the different concentrations of ammonia, nitrite and nitrate are shown in Figure 1.




Figure 1. Growth of juvenile *Farfantepenaeus brasiliensis* at different concentrations of ammonia (A), nitrite (B) and nitrate (C) over 40 days. Bars = standard error.

The survival rates of *F. brasiliensis* juveniles exposed to ammonia, nitrite and nitrate during

the experiment are expressed in Figure 2. No significant differences were observed ($p > 0.05$)

among treatments TAN-50%, TAN-SL and the control from day 0 to 30. However, at the end of the experiment, treatments TAN-50% and TAN-SL differed significantly ($p < 0.05$) from the control group (Figure 2a).

Among treatments NO_2 -50%, NO_2 -SL and the control, there were no significant differences ($p > 0.05$) on days 0, 10 and 30. On day 20, there were significant differences ($p < 0.05$) between the control and groups NO_2 -50% and NO_2 -SL.

After 40 days, all of the groups differed significantly (Figure 2b).

On days 0 and 10, no significant differences were observed ($p > 0.05$) between treatments NO_3 -50%, NO_3 -SL and the control. Significant differences ($p < 0.05$) between the control and groups NO_3 -50% and NO_3 -SL appeared on day 20; after 40 days, there were significant differences among all groups (Figure 2c).




Figure 2. Mean survival values of *Farfantepenaeus brasiliensis* juveniles exposed to different concentrations of ammonia (A), nitrite (B) and nitrate (C) over 40 days.

DISCUSSION

The water temperature during the experiment remained within the adequate range of 24 to 32 °C for the growth of penaeid shrimp (VAN WYK and SCARPA, 1999). The true safe levels for different species of farmed aquatic organisms may differ markedly from those obtained in short-term tests (TOMASSO, 1994). The safe levels proposed by this method, used by CAMPOS *et al.* (2012), can be subject to errors in the estimation of the concentrations that are effectively chronic to *F. brasiliensis*, as evidenced by the results of this study.

High concentrations of nitrogen compounds (ammonia, nitrite and nitrate) affect physiological processes that are important to aquaculture activities (MONTTOYA *et al.*, 1999). Among these processes, affected osmoregulation and respiration result in low food consumption, low specific growth rates and even mortality of the shrimp (WASIELESKY, 2000; KUHN *et al.*, 2010, BARBIERI 2010; CAMPOS *et al.*, 2012; BARBIERI *et al.*, 2014). During the period of molting, shrimp can be more sensitive to the different effects of the nitrogenous compounds, such as ammonia, nitrite

and nitrate, which are also more toxic during this period (WASIELESKY, 2000).

Ammonia is the end product of protein catabolism for most aquatic organisms (KINNE, 1976) and is present in the aquatic environment in the ionized (NH_4^+) and non-ionized form (NH_3), which can spread through cell membranes. The process of ammonia ionization is regulated by the pH level such that the percentage of NH_3 increases and the percentage of NH_4^+ decreases in relation to increasing pH levels (FROMM and GILLETTE, 1968). In this context, after a 30-day exposure to ammonia, the survival of the shrimp in the treatments TAN-50% and TAN-SL did not show significant differences in comparison with the control group. However, during this period, the growth in wet weight demonstrated that a concentration as low as the safe level, as determined by CAMPOS *et al.* (2012), caused growth retardation.

High levels of ammonia can be harmful to crustaceans and cause various adverse effects such as reduced growth and survival (LIN and CHEN, 2001; BARBIERI, 2010; COBO *et al.*, 2014). WASIELESKY *et al.* (1994) confirmed that the growth rates of *Farfantepenaeus paulensis* (post-larvae) were significantly reduced in ammonia concentrations of 0.07 to 0.14 mg L^{-1} . MIRANDA-FILHO *et al.* (2009) studied juveniles of the same species in pre-nursery and nursery stages for 75 days and observed reduced predation activity and growth.

Nitrite is the intermediate compound in the bacterial nitrification of ammonia into nitrate (in oxidizing environments) or the production of nitrate denitrification (in reducing environments) and is toxic to aquatic organisms, causing mortality in culture systems (BROWNELL, 1980; TSAI and CHEN, 2002). The ionization process of nitrite is also mediated by pH levels, so that the percentage of nitrous acid (HNO_2) increases with decreasing pH levels. Introduction of high concentrations of nitrite into the aquatic environment may lead to hemolymphatic problems because its toxic action targets the process of oxygen transport, converting hemocyanin into metahemocyanin, which is unable to carry oxygen to the tissues (GROSS, 2004). This process decreases the amount of

oxygen available for metabolism (TAHON *et al.*, 1988) and leads to hypoxia and significant mortality (CHEN *et al.*, 1986; BARBIERI *et al.*, 2014).

The results of the present study demonstrated that juveniles of *F. brasiliensis* only showed significant reduction in weight gain after 30 days of exposure. However, after 20 days of exposure to nitrite, the survival of the shrimp in treatments NO_2 -50% and NO_2 -SL showed significant differences in relation to the control group. GROSS (2004) reported growth retardation and mortality in *L. vannamei* in farms located in Israel, which could be attributed to the high concentrations of nitrite (8 mg L^{-1}). CHEN and CHEN (1992) exposed juvenile *Penaeus monodon* to nitrite concentrations that ranged between 2 and 20 mg L^{-1} for 60 days. The shrimp exposed to 4, 8 and 20 mg L^{-1} of nitrite showed significantly lower weight gain. WASIELESKY (2000) studied juvenile *F. paulensis* and verified a significant reduction in weight gain at a concentration of 20.4 $\text{mg NO}_2 \text{ L}^{-1}$ (equivalent to twice the safe level) after the 30-day trial. In addition, this species showed a mortality rate of 61% at a nitrite concentration of 10.2 mg L^{-1} . However, in spite of the high mortality rate, the growth of the shrimp that survived was not significantly affected by nitrite. According to the author, this fact is likely related to the greater requirement for oxygen during the process of molting. This suggests that shrimp that manage to pass through the molting stage (ecdysis) have relatively normal weight gain for a period of time even under these elevated nitrite concentrations.

In systems of culture LIN and CHEN (2003) verified that nitrite concentrations equal to or higher than 25.7 mg L^{-1} and a salinity of 35 can reduce the growth of *L. vannamei*. FURTADO *et al.* (2011) did not observe significant effects of nitrite on *L. vannamei* in the treatments with 3.1 and 4.3 mg L^{-1} . While MAICÁ *et al.* (2011) worked with different salinities in the bioflocs system and verified that the highest nitrite levels occurred at salinities of 2 and 25. They also observed that there was mortality of the shrimp when the nitrite concentrations exceeded the safe levels for *L. vannamei* juveniles ($\leq 1 \text{ mg L}^{-1}$) (VAN WYK and SCARPA, 1999) between 20 and 30 days. The highest mortality rates and the highest nitrite concentrations that occurred at a salinity of 2

suggest that the nitrite level is one of the main factors that cause mortality. BARBIERI *et al.* (2014) showed the inverse relationship between salinity and nitrite toxicity for juveniles of *Litopenaeus schmitti* and emphasized that the toxicity increases when animals are exposed to a hyposmotic conditions.

Nitrate is the least toxic nitrogen compound for aquatic organisms including Penaeidae (VAN WYK and SCARPA, 1999), although its study is important because it can produce lethal or sublethal effects on different organisms or act synergistically with other nitrogenous substances (KUHN *et al.*, 2009). KUHN *et al.* (2010) showed that *L. vannamei* can be cultivated under a salinity of 11 with 220 ppm of nitrate for a period of six weeks, while nitrate levels of 435 ppm were not safe for these shrimp. The growth rates of the shrimp *F. paulensis* exposed to 80.7 mg L⁻¹ of nitrate were significantly lower in relation to the control treatment (WASIELESKY, 2000). In the present study, the growth and survival of *F. brasiliensis* juveniles exposed for 40 days to concentrations of 45.6 and 91.2 mg L⁻¹ of nitrate were negatively affected in comparison with the control group. In this context, this study confirms the results of the authors mentioned above and emphasizes the greater sensitivity of *F. brasiliensis* to high levels (90 mg L⁻¹) of nitrate compared with that of the white shrimp *L. vannamei*.

According to FRIAS-ESPERICUETA *et al.* (1999), the interaction between nitrogenous compounds and shrimp production is an important consideration for farmers. There is no relevant accumulation of nitrogenous compounds in semi-intensive farming systems with low stocking densities and high rates of water renewal. Conversely, super-intensive systems with biofilter or bioflocs technology and minimal water renewal rates feature a natural process of nitrate accumulation in the system (> 100 mg L⁻¹) (POERSCH *et al.*, 2007, FURTADO *et al.*, 2011). SOUZA *et al.* (2014) verified ammonia and nitrite levels of 6.0 and 10 mg L⁻¹, respectively, at the stage of biofloc formation in the culture of *F. brasiliensis*. EMERENCIANO *et al.* (2012) found that juvenile *F. brasiliensis* reared in a bio-flocs technology (BFT) system showed higher final weight and weight gain than juveniles cultivated in clear water and highlighted the potential of this species

in super-intensive systems with minimal water renewal.

CONCLUSIONS

The results of this study emphasize the sensitivity of this species to nitrate as the worst survival results were shown in the treatment at the supposed safe level. Thus, special care should be taken to control the concentrations of nitrogen products in *F. brasiliensis* farming because they can significantly affect the final output of this species in aquaculture systems.

ACKNOWLEDGMENTS

The authors are grateful for the financial support provided by the National Council for Scientific and Technological Development (CNPq), Ministry of Fishery and Aquaculture (MPA) and Coordination for the Improvement of Higher Level Personnel (CAPES). W.J. Wasielesky, F. D'Incao and L.H. Poersch are research fellows of CNPq. Bruno Campos received the CAPES grants.

REFERENCES

- AMINOT, A. e CHAUSSEPIED, M. 1983 *Manuel des analyses chimiques en milieu marin*. Centre National pour l'Exploitation des Océans (CNEXO), Brest, France. 395p.
- BARBIERI, E. 2010 Acute toxicity of ammonia in white shrimp (*Litopenaeus schmitti*) (Burkenroad, 1936, Crustacea) at different salinity levels. *Aquaculture*, 306(1-4): 329-333.
- BARBIERI, E.; BONDIOLI, A.C.V.; MELO, C.B.; HENRIQUES, M.B. 2014 Nitrite toxicity to *Litopenaeus schmitti* (Burkenroad, 1936, Crustacea) at different salinity levels. *Aquaculture Research*, 1-9. doi: 10.1111/are.12583.
- BENDSCHNEIDER, K. e ROBINSON, R.J. 1952 A new spectrophotometric method for the determination of nitrite in sea water. *Journal of Marine Research*, 11: 87-96.
- BROWNELL, C.L. 1980 Water quality requirements for first feeding in marine fish larvae: ammonia, nitrite and nitrate. *Journal of Experimental Marine Biology and Ecology*, 44(2): 269-283.

- CAMPOS, B.R.; MIRANDA-FILHO, K.; D'INCAO, F.; POERSCH, L.; WASIELESKY, W. 2012 Toxicidade aguda da amônia, nitrito e nitrato sobre os juvenis de camarão-rosa *Farfantepenaeus brasiliensis* (Latreille, 1817) (Crustacea: Decapoda). *Atlântica*, 34(1): 75-81.
- CHEN, J.C. and CHEN, S.F. 1992 Effects of nitrite on growth and molting of *Penaeus monodon* juveniles. *Comparative Biochemistry and Physiology C*, 101(3): 453-458.
- CHEN, J.C.; CHIN, C.K.; LEE, C.K. 1986 Effects of ammonia and nitrite on larval development of the shrimp *Penaeus monodon*. *The first Asian fisheries forum. Asian fishery Society*. p.657-662.
- COBO, M.L.; SONNENHOLZNER, S.; WILLE, M.; SORGELOOS, P. 2014. Ammonia tolerance of *Litopenaeus vannamei* (Boone) larvae. *Aquaculture Research*, 45(3): 470-475.
- EMERENCIANO, M.; BALLESTER, E.L.C.; CAVALLI, R.O.; WASIELESKY, W. 2012 Biofloc technology application as a food source in a limited water exchange nursery system for pink shrimp *Farfantepenaeus brasiliensis* (Latreille, 1817). *Aquaculture Research*, 43(3): 447-457.
- FRIAS-ESPERICUETA, M.G.; HARFUSH-MELENDEZ, M.; OSUNA-LSPEZ, J.I.; PAEZ-OSUNA, F. 1999 Acute toxicity of ammonia to juvenile shrimp *Penaeus vannamei* Boone. *Bulletin of Environmental Contamination and Toxicology*, 62(5): 646-652.
- FROMM, P.O. and GILLETE, J.R. 1968 Effect of ambient ammonia on blood ammonia and nitrogen excretion of rainbow trout (*Salmo gairdneri*). *Comparative Biochemistry and Physiology*, 26(3): 887-896.
- FURTADO, P.S.; POERSCH, L.H.; WASIELESKY, W. 2011 Effect of calcium hydroxide, carbonate and sodium bicarbonate on water quality and zootechnical performance of shrimp *Litopenaeus vannamei* reared in bio-flocs technology (BFT) systems. *Aquaculture*, 321(1-2): 130-135.
- GROSS, A. 2004 Acute and chronic effects of nitrite on white shrimp, *Litopenaeus vannamei*, cultured in low-salinity brackish water. *Journal of the World Aquaculture Society*, 35(3): 315-321.
- JENSEN, L.V. 2012 *Produção e transporte do camarão-rosa Farfantepenaeus brasiliensis para a pesca amadora*. São Carlos, 133p. (Tese de Doutorado. Universidade Federal de São Carlos). Available at: <http://www.bdt.ufscar.br/htdocs/tedeSimplificado//tde_busca/arquivo.php?codArquivo=5407>
- KINNE, O. 1976 Cultivation of marine organisms: water quality management of technology. In: KINNE, O. *Marine Ecology*. New York: Wiley interscience. Vol III, part 1. p.79-300.
- KUHN, D.D.; BOARDMAN, G.D.; MARSH, L.; LAWRENCE, A.L.; FLICK, G.J. 2009 Technology and research advances for the production of marine shrimp in recirculating aquaculture systems. *Journal of Shellfish Research*, 28: 709p.
- KUHN, D.D.; SMITH, S.A.; BOARDMAN, G.D.; ANGIER, M.W.; MARSH, L.F.J.; GEORGE, J. 2010 Chronic toxicity of nitrate to Pacific white shrimp, *Litopenaeus vannamei*: Impacts on survival, growth, antennae length, and pathology. *Aquaculture*, 309(1-4): 109-114.
- LIN, Y.C. and CHEN, J.C. 2001 Acute toxicity of ammonia on *Litopenaeus vannamei* Boone juveniles at different salinity levels. *Journal of Experimental Marine Biology and Ecology*, 259(1): 109-119.
- LIN, Y.C. and CHEN, J.C. 2003 Acute toxicity of nitrite on *Litopenaeus vannamei* (Boone) juveniles at different salinity levels. *Aquaculture*, 224(1-4): 193-201.
- LOPES, D.A.; PEIXOTO, S.R.M.; WASIELESKY, W.; BALLESTER, E.L.C. 2009 Análise comparativa da criação dos camarões-rosa *Farfantepenaeus brasiliensis* e *Farfantepenaeus paulensis* criados em gaiolas em ambiente estuarino. *Ciência Rural*, 39(9): 1540-1546.
- MAICÁ, P.F.; BORBA, M.R.; WASIELESKY, W. 2011 Effect of low salinity on microbial floc composition and performance of *Litopenaeus vannamei* (Boone) juveniles reared in a zero-water-exchange super-intensive system. *Aquaculture Research*, 43(7): 361-370.
- MIRANDA-FILHO, K.C.; PINHO, G.L.L.; WASIELESKY, W.; BIANCHINI, A. 2009 Long-term ammonia toxicity to the pink-shrimp *Farfantepenaeus paulensis*. *Comparative Biochemistry and Physiology C*, 150(3): 377-382.
- MONTOYA, R.A.; LAWRENCE, A.L.; GRANT, W.E.; VELASO, M. 1999. Simulation of nitrogen

- dynamics and shrimp growth in an intensive shrimp culture system: effects of feed and feeding parameters. *Ecological Modelling*, 122(1-2): 81-95.
- OSTRENSKY, A. and WASIELESKY, W. 1995 Acute toxicity of ammonia to various life stages of the São Paulo shrimp *Penaeus paulensis* Pérez-Farfante, 1967. *Aquaculture*, 132(3-4): 339-347.
- POERSCH, L.H.; SANTOS, M.H.S.; MIRANDA, K.F.; WASIELESKY, W. 2007 Efeito agudo do nitrato sobre alevinos da tainha *Mugil platanus* (Pisces: Mugilidae). *Boletim do Instituto de Pesca*, 33(2): 247-252.
- RAND, G.M. and PETROCELLI, P.R. 1985 *Fundamentals of aquatic toxicology*. 1º ed. Taylor & Francis, USA. 666p.
- SOUZA, D.M.; SUITA, S.M.; ROMANO, L.A.; WASIELESKY, W.Jr.; BALLESTER, E.L.C. 2014 Use of molasses as a carbon source during the nursery rearing of *Farfantepenaeus brasiliensis* (Latreille, 1817) in a biofloc technology system. *Aquaculture Research*, 45(2): 270-277.
- TAHON, J.P.; VAN HOOFF, D.; VINCKIER, C.; WITTERS, R.; DE LEY, M.; LONHE, R. 1988 The reaction of nitrite with the haemocyanin of *Astacus leptodactylus*. *Biochemical Journal*, 249(3): 233-242.
- TSAI, S.J. e CHEN, J.C. 2002 Acute toxicity of nitrate on *Penaeus monodon* juveniles at different salinity levels. *Aquaculture*, 213(1-4): 163-170.
- TOMASSO, J.R. 1994 Toxicity of nitrogenous Wastes to Aquaculture Animals. *Reviews in Fisheries Science*, 2(4): 291-314.
- UNESCO. 1983 *Chemical methods for use in marine environmental monitoring. Manual and Guides 12*. Intergovernmental Oceanographic Commission. Paris, France. 53p.
- VAN WYK, P. and SCARPA, J. 1999 Water quality requirements and management. In: VAN WYK, P. *Farming Marine Shrimp in Recirculating Freshwater Systems*. Tallahassee: Florida Department of Agriculture and Consumer Services. p.128-138.
- WAJSBROT, N.; GASITH, A.; DIAMANT, A.; POPPER, D.M. 1993 Chronic toxicity of ammonia to juvenile gilthead seabream *Sparus aurata* and related histopathological effects. *Journal of Fish Biology*, 42(3): 321-328.
- WASIELESKY, W.J. 2000 *Cultivo de juvenis do camarão-rosa Farfantepenaeus paulensis* (Decapoda, Penaeidae) no estuário da Lagoa dos Patos: efeitos dos parâmetros ambientais. Rio Grande. 147p. (Tese de Doutorado. Universidade Federal de Rio Grande).
- WASIELESKY, W.J.; MARCHIORI, M.A. (in memorian); SANTOS, M.H.S. 1994. Efeito da amônia no crescimento de pós-larvas do camarão rosa, *Penaeus paulensis*, Pérez-Farfante, 1967 (Decapoda:Penaeidae). *Nauplius*, 2: 99-105.
- WASIELESKY, W.; PEIXOTO, S.; JENSEN, L.; POERSCH, L.H.; BIANCHINI, A. 2004 Estudo preliminar do cultivo do camarão-rosa *Farfantepenaeus paulensis* em cercados no estuário da Lagoa dos Patos. *Boletim do Instituto de Pesca*, 30(1): 63-70.
- ZAR, J.H. 1996 *Biostatistical analysis*. 3.ed. New Jersey: Prentice Hall. 662p.